Tel: 22003434

Website: www.sth.org.hk

- 本院设有「八达通」时钟泊车服务 We provide Octopus hourly parking service
- 公共小巴绿色专线/ Green Line Public Bus: 2, 17M, 25M, 46, 70
- 红色非专线: 旺角往新蒲岗/黄大仙/九龙 城

Red Line Minicab: Mongkok to San Po Kong / Wong Tai Sin/Kowloon City

- 公共巴士/ Public Bus: 1, 1A, 2A, 6D, 7B, 9, 12A, 13D, 16, 24, 27, 42, 95, 98C, 113, 203E, 296C, N216
- 港铁 / MTR
- 乐富站换乘的士约五分钟到达 / Lok Fu Station: 5 mins by taxi
- 旺角站换乘非专线小巴(近先达广场)至露 明道 Mongkok Station: by red line minicab, get off at Lomond Road
- 宋皇台: B1 出口, 可步行 5-10 分到达 Sung Wong Toi: Exit B1, about 5-10 minutes walk
- 东铁 / East Rail Line
- 旺东可换乘的士约五分钟到达 Mongkok East:about 5 minutes by taxi
- * 九龙塘可换乘专线小巴 25M 至界限街 Kowloon Tong:by green line bus no. 25M
- 复康巴士电召服务/Rehabus (Dial a Ride): 28178154

Effective Date: September 2022

PFE-DMC-2-S010





St. Teresa's Hospital 聖德肋撒醫院

Management of Diarrhea 腹泻处理

Health Information 健康信息

腹泻处理

- 多喝开水或稀粥水
- 勿进食油腻煎炸、高纤维、 蔬果类及奶类食
- 进食前及如厕后必须洗手
- 按医生处方服用补充水份溶液
- 如腹泻加剧、小便量减少、 呼吸加速、烦躁不安、神志 不清或呆滞,请尽快求医
- 替小孩患者更换尿片后, 必须替其清洗,臀部可涂 上合适药膏以保护臀部皮 肤

*以上信息只供参考,并不能完全反映医生意见,想了解更多有关信息,应咨询你的医生。

Management of Diarrhea

- Drink more water or congee
- DO NOT take fried food, food containing high fiber, vegetables or dairy products
- Take oral re-hydrating solution as doctor's prescribed



- Seek medical help immediately if:
- diarrhea gets more severe
- decreased urine output
- breathing getting faster
- # found restless
- found unconscious or delirious
- For patients who are incontinent, please rinse the buttock whenever napkin is changed.
 Suitable lotion can be applied to prevent skin laceration on buttock

References: World Health Organization. (2007). Guideline for management of common diseases in young children in emergencies

Revised Date: 1 September 2013

^{*}This document is for information purpose and is not intended to be a substitute for the advice of a doctor. Should you have any queries, please consult your doctor-incharge.