

Side Effects of Chemotherapy

The most common side effects are bone marrow suppression, such as decreased in red blood cell, white blood cell or platelet count. Patient has the potential to get infection and feel tired. Other side effects include nausea, vomiting, alopecia, constipation, diarrhea, sore mouth, muscle pain, tingling sensation in hands or feet and infertility.



Side Effects of Targeted Therapy

They include diarrhea, rash, itching, dry or cracked skin, acne, nail changes, mouth infections and Pneumonitis.



Side Effects of Immunotherapy

Common side effects are similar to targeted therapy.

Rare side effects include:

Immune-Mediated Pneumonitis, Colitis, Hepatitis, Nephritis and Endocrinopathies, e.g. Thyroid dysfunction..



Tel: (852) 22003434

Website: www.sth.org.hk


- We provide "Octopus hourly parking services"
- Green Line Minicab: 2, 17M, 25M, 46, 70,
- Red Line Minicab: Mongkok to San Po Kong / Wong Tai Sin / Kowloon City
- Public Buses: 1, 1A, 2A, 6D, 7B, 9, 12A, 13D, 16, 24, 27, 42, 95, 98C, 113, 203E, 296C, N216
- MTR
- * Lok Fu: 5 mins by taxi
- * Mongkok: by red line minibus, get off at Lomond Road
- * Sung Wong Toi: Exit B1, about 5-10 minutes walk
- East Rail Line
- * Mongkok East: about 5 minutes by taxi
- * Kowloon Tong: by green line bus no. 25M
- Rehabus (Dial-a Ride): (852) 2817 8154



St. Teresa's Hospital
聖德肋撒醫院

Oncology Centre

Anti-Cancer
Intravenous Therapy



Anti-Cancer Intravenous Therapy

Chemotherapy

It is a kind of drug treatment which is used to kill cancer cells or prevent their growth in order to reduce signs and symptoms.




Targeted Therapy

It works by blocking and interfering the signal transmission of cells in order to control the growth of cancer cells. Besides, it minimizes the damage to normal cells during the treatment.



Immunotherapy

It uses certain parts of a person's immune system to fight cancer. It stimulates your own immune system to work harder or smarter to attack cancer cells.



How Do Anti-Cancer Drugs Work?

Anti-cancer drugs used based on the stage of disease, the histopathology or genetic study report. Anti-cancer drugs are commonly administered via intravenous (IV), (SC), or oral route. No matter which route is used, the drug can enter the body and attack cancerous cells.



Method of Intravenous injection

Most drugs are given by intravenous infusion after diluted as prescribed by doctor. It is administered from a few minutes to a few hours (out-patient) or a few days (in-patient) depending on an individual's regimen.



Phone: (852) 22003193
Fax: (852) 27116747
Website: www.sth.org.hk
Email: oncology@sth.org.hk

Procedures before IV Therapy

1. Patient, relatives will be arranged an appointment to consult Oncologist before starting the treatment. The Oncologist explains the effect, side effect and the regimen of the IV therapy to the patient.
2. Patient's significant others and relatives are encouraged to accompany each consultation and treatment.
3. Blood or urine test before each therapy.
4. Informed consent should be signed before treatment by patient / significant others.
5. The dosage of anti-cancer drugs will be prescribed by the oncologist according to general condition, blood result, body weight and body height.
6. Patient / relatives will go to the Pharmacy to pay for the medication and bring back to Oncology Centre. Our staff will collect the chemo drug from pharmacy after the reconstitution in pharmacy.
7. IV therapy will be started after the drug is ready.
8. Patient's vital signs will be monitored & recorded before and after each therapy for evaluation of his / her condition.



Follow up

Patient will be arranged Investigation and follow up appointment after each therapy.

If patient feel discomfort / sick after the therapy, he /she can contact us within office hours for earlier appointment or directly contact oncologists in-charge after office hours.

(This leaflet is for reference only)

